

Anomalous t - c - g coupling: The Connection between Single Top Production and Top Decay

Tim Tait and C.-P. Yuan

Department of Physics and Astronomy, Michigan State University
East Lansing, MI 48824, USA

Abstract

Continuing earlier work, we examine the constraint on an anomalous t - c - g coupling from top quark decay. We find that from current CDF measurements of the branching ratio $t \rightarrow Wb$, the minimum scale at which new physics can strongly modify the t - c - g coupling is $\Lambda_{tcg} \geq$ about 950 GeV. At the upgraded Tevatron, single top production can constrain $\Lambda_{tcg} \geq 4.5$ TeV. The connection between t - c production and the decay $t \rightarrow cg$ is examined, showing how constraints on one lead to a constraint on the other.

As shown in [1], an anomalous coupling between top quark, charm quark, and gluon fields can have a large effect on single top production at the Tevatron¹. In that work, it was mentioned that the t - c - g operator can also be studied from the decay $t \rightarrow cg$. This decay mode has been studied in [2], and the results are similar to those presented here.

In [1] the ratio of the partial widths is presented,

$$R_{tcg} = \frac{\Gamma(t \rightarrow cg)}{\Gamma(t \rightarrow W^+b)} = \frac{\sqrt{2} \, 64\alpha_s \pi m_t^2}{3\Lambda^2 G_F \left[1 - \frac{m_W^2}{m_t^2}\right]^2 \left[1 + 2\frac{m_W^2}{m_t^2}\right]}. \quad (1)$$

¹ The notation and conventions used in this work are the same as those presented in [1]. A full discussion of the effective operator which generates the t - c - g coupling can be found there-in.

Assuming that there is no other new physics modifying the top quark decays, the branching ratio $\text{BR}(t \rightarrow Wb)$ can be expressed as,

$$\text{BR}(t \rightarrow Wb) = \frac{\Gamma(t \rightarrow Wb)}{\Gamma(t \rightarrow Wb) + \Gamma(t \rightarrow cg)}. \quad (2)$$

From these results, one may determine the Λ_{tcg} which corresponds to a given lower limit on $\text{BR}(t \rightarrow Wb)$, b_{min} . The results are plotted in Figure-1.




Figure 1: The constraint on Λ_{tcg} as a function of b_{min} .

The current CDF measurement of $\text{BR}(t \rightarrow Wb)$,

$$\text{BR}(t \rightarrow Wb) = 0.87^{+0.13 \pm 0.13}_{\pm 0.30 \pm 0.11} \quad (3)$$

leads to the 1σ limit $b_{min} = 0.55$. From this we derive a constraint on $\Lambda_{tcg} \geq 950$ GeV. As shown in [1], the maximum effect to the top-charm production

rate (99% C.L.) is,

$$\sigma_{tc}^{max} = 616 \left(\frac{\text{TeV}}{\Lambda_{tcg}} \right)^2 \text{ fb}, \quad (4)$$

including some basic acceptance cuts. From the constraint on Λ_{tcg} discussed above, this implies $\sigma_{tc}^{max} \leq$ about 0.6 pb. The relation between b_{min} and σ_{tc}^{max} is plotted in Figure-2. At the upgraded Tevatron (Run-II), the integrated luminosity is expected to be 2 fb^{-1} per year. As shown in [1], if no t - c signal events are observed in 2 fb^{-1} of integrated luminosity, this would imply $\Lambda_{tcg} \geq 4.5 \text{ TeV}$, and from Figure-1, this requires $b_{min} = 96.3\%$ ($\text{BR}(t \rightarrow cg) \leq 3.7\%$).




Figure 2: The maximum cross section, σ_{tc}^{max} as a function of b_{min} .

The connection between σ_{tc}^{max} and b_{min} is interesting, because it gives a prediction of one quantity in terms of the other one. This can serve to help

identify this type of new physics effect from other possible effects, and shows how a constraint derived from one process has implications for another one.

Recently detailed work has been done on studying the measurement of Λ_{tcg} from top decays, including kinematic cuts to optimize the signal to background ratio [3]. These results show studying top decays to be a promising way to constrain Λ_{tcg} , almost as promising as studying single-top production for the current data sample, and possibly providing even stronger constraints at the upgraded Tevatron.

The authors would like to thank Ehab Malkawi and X. Zhang for helpful discussion. This work was supported in part by the NSF under grant numbers PHY-9309902 and PHY-9507683.

References

- [1] E. Malkawi and T. Tait, Michigan State preprint MSUHEP-51116 (1995).
- [2] T. Han, K. Whisnant, B.-L. Young, and X. Zhang, hep-ph/9603247, March 1996.
- [3] T. Han, K. Whisnant, B.-L. Young, X. Zhang, hep-ph/9606231, May 1996.

- [4] J. Incandela (CDF Collaboration), FERMILAB-CONF-95-237-E (July 1995).